

## Could Mindful Meditation be a Routine Therapeutic Intervention in Practice?

Jones Gupta\*

University of Alberta, 116 St & 85 Ave, Edmonton, AB T6G 2R3, Canada

\*: All correspondence should be sent to: Dr. Jones Gupta

Author's Contact: Dr. Jones Gupta, M.D., Ph.D., E-mail: [jones.gupta.ca@gmail.com](mailto:jones.gupta.ca@gmail.com)

DOI: <https://doi.org/10.15354/si.24.co202>

Funding: No funding source declared.

COI: The author declares no competing interest.

AI Declaration: The author affirms that artificial intelligence did not contribute to the process of preparing the work.

Although mindfulness meditation has become more and more well-liked as a therapeutic technique in recent years, it might not be appropriate for every person or situation. Although mindfulness meditation has been shown to be beneficial for reducing stress, anxiety, and depression, its effects may not be felt by all people. Furthermore, some people could find it challenging to practice mindful meditation because of a variety of obstacles, like attentional or cognitive impairments. It is pivotal to consider the unique demands of every client in a professional context and design interventions accordingly. It's also critical to understand that, where appropriate, mindful meditation should be utilized in addition to other evidence-based therapies rather than being viewed as a one-size-fits-all remedy. Because of this, mindful meditation should not be used as the only therapeutic intervention in practice, even if it can be a useful tool in increasing mental well-being.

**Keywords:** Mindful Meditation; Clinical Practice; Benefits; Risks; Therapeutic

Science Insights, 2024 July 31; Vol. 45, No. 1, pp.1421-1423.

© 2024 Insights Publisher. All rights reserved.



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the [Creative Commons Attribution-NonCommercial 4.0 License](https://creativecommons.org/licenses/by-nc/4.0/) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed by the Insights Publisher.

**M**INDFUL meditation has become increasingly popular as a potential therapeutic strategy for a range of mental health issues. Mindfulness entails directing one's attention to the current moment and embracing one's thoughts and emotions without evaluation (1). One of the primary advantages of practicing mindful meditation is its capacity to assist people in effectively coping with stress and worry (2). Through the practice of training, one's mind concentrates on the current moment, individuals might acquire the ability to release troubling ideas and emotions that contribute to feelings of ten-

sion and anxiety. This can result in an enhanced state of tranquility and serenity, which can exert a beneficial influence on one's mental well-being.

Mindful meditation can serve as a valuable resource for persons suffering with depression. Mindful meditation promotes the acceptance of one's thoughts and feelings without judgment, which can lead to the cultivation of enhanced self-compassion and self-acceptance (3). This can be particularly advantageous for people who experience difficulties with low self-esteem and self-criticism, which are prevalent in cases of depression. Fur-

thermore, mindful meditation has demonstrated efficacy in enhancing general well-being, in addition to its positive effects on stress, anxiety, and depression. Studies have discovered that consistent engagement in mindful meditation can result in heightened self-awareness, improved attention and concentration, and an enhanced sense of inner tranquility (4, 5). These advantages can significantly influence persons' quality of life and general mental well-being.

Incorporating mindful meditation into established therapy procedures can be highly beneficial. Therapists can assist individuals in cultivating enhanced emotional regulation and resilience by instructing them on how to be fully present in the present moment and embrace their thoughts and sensations without passing judgment. This is particularly crucial for people who grapple with overwhelming emotions or encounter difficulties in managing the obstacles of life. Mindful meditation can serve as a significant resource for persons who have experienced trauma or have been diagnosed with posttraumatic stress disorder (PTSD) (6). Mindful meditation can assist individuals in cultivating heightened self-awareness and self-compassion, facilitating the processing of traumatic experiences and alleviating symptoms associated with PTSD. This can serve as a beneficial supplement to conventional therapeutic methods for trauma, such as cognitive-behavioral therapy or exposure therapy. In addition, practicing mindful meditation can be a cost-effective and easily accessible kind of therapy (7). Contrary to conventional therapy methods that often necessitate numerous appointments with a therapist, mindful meditation can be performed autonomously and at one's convenience. This feature might be a great resource for persons who lack access to conventional therapy or who desire supplementary support beyond their therapy sessions.

Nevertheless, while certain individuals may find it beneficial in managing their stress and anxiety, others may not derive the same advantages. This is due to the fact that every individual possesses distinct characteristics, and strategies that are effective for one person may not yield the same results for another. Hence, it is crucial to contemplate alternative therapeutic methods that may be better suited for individuals who do not derive benefits from mindful meditation (8). In addition, successful mindful meditation demands a significant amount of time and unwavering commitment. In the contemporary day characterized by rapidity, not all individuals possess the privilege of allocating time for the habitual engagement in meditation. This can be a challenge for patients seeking to integrate mindful meditation into

their daily schedule, perhaps diminishing its efficacy as a therapeutic intervention (9). Moreover, certain people may encounter difficulty in maintaining concentration and mindfulness when engaging in meditation, hence diminishing its efficacy as a therapeutic tool.

Another factor that prevents mindful meditation from being a routine therapeutic practice is its potential inability to effectively target the root causes of mental health problems. Although meditation can assist individuals in effectively managing their symptoms and enhancing their general state of being, it may not directly target the underlying source of their issues (10). To achieve genuine healing and conquer mental health challenges, individuals may need to delve into alternative therapeutic approaches that specifically target their unique needs and problems. Further, it is important to note that mindful meditation may not be appropriate for those with specific mental health disorders. Individuals with severe depression or anxiety may encounter difficulties in engaging in meditation due to the presence of their symptoms (11). In such instances, alternative therapeutic methods such as cognitive-behavioral therapy or medication may prove more efficacious in addressing their mental health concerns. So, it is crucial to consider the individual's mental health background and requirements while deciding on the most suitable therapeutic intervention. In addition, mindful meditation may not be appropriate for persons who have a history of trauma or complicated mental health conditions (12). For these individuals, alternative therapeutic methods that target their specific needs and concerns may be more helpful in facilitating recovery and enhancing overall well-being. When deciding on the most suitable therapeutic intervention for an individual, it is critical to consider their specific circumstances and personal history.

Moreover, persons who struggle with introspection or self-reflection may find mindful meditation unsuitable (13, 14). These individuals may encounter difficulties while attempting to delve into their thoughts and emotions during meditation, which can restrict the efficacy of this practice as a therapeutic tool. In such instances, alternative therapy methods that prioritize the development of self-awareness and insight may yield greater benefits. As thus, mindful meditation is not a universally applicable method of therapy. While meditation may be advantageous for many individuals, others may get better results with alternative therapeutic approaches (15). It is crucial for patients to examine several alternatives and determine the most effective method for managing their mental health and well-being. ■

## References

1. Schuman-Olivier Z, Trombka M, Lovas DA, Brewer JA, Vago DR, Gawande R, Dunne JP, Lazar SW, Loucks EB, Fulwiler C. Mindfulness and behavior change. *Harv Rev Psychiatry* 2020; 28(6):371-394. DOI: <https://doi.org/10.1097/HRP.0000000000000277>
2. Bostock S, Crosswell AD, Prather AA, Steptoe A. Mindfulness on-the-go: Effects of a mindfulness meditation app on work stress and well-being. *J Occup Health Psychol* 2019; 24(1):127-138. DOI:

<https://doi.org/10.1037/ocp0000118>

3. Centeno RPR. Effect of mindfulness on empathy and self-compassion: An adapted MBCT Program on Filipino college students. *Behav Sci (Basel)* 2020; 10(3):61. DOI: <https://doi.org/10.3390/bs10030061>
4. Ho CYY, Yu BCL, Mak WWS. Nonattachment mediates the associations between mindfulness, well-being, and psychological distress: A meta-analytic structural equation modeling approach. *Clin Psychol Rev* 2022; 95:102175. DOI: <https://doi.org/10.1016/j.cpr.2022.102175>
5. Chan RMK, Mak WWS, Yu BCL. Going beyond mindfulness: How concentration and tranquility commonly co-arising with mindfulness account for mental health. *Int J Environ Res Public Health* 2023; 20(8):5470. DOI: <https://doi.org/10.3390/ijerph20085470>
6. Boyd JE, Lanius RA, McKinnon MC. Mindfulness-based treatments for posttraumatic stress disorder: A review of the treatment literature and neurobiological evidence. *J Psychiatry Neurosci* 2018; 43(1):7-25. DOI: <https://doi.org/10.1503/jpn.170021>
7. Behan C. The benefits of meditation and mindfulness practices during times of crisis such as COVID-19. *Ir J Psychol Med* 2020; 37(4):256-258. DOI: <https://doi.org/10.1017/ipm.2020.38>
8. Rich T, Chrisinger BW, Kaimal R, Winter SJ, Hedlin H, Min Y, Zhao X, Zhu S, You SL, Sun CA, Lin JT, Hsing AW, Heaney C. Contemplative practices behavior is positively associated with well-being in three global multi-regional Stanford WELL for life cohorts. *Int J Environ Res Public Health* 2022; 19(20):13485. DOI: <https://doi.org/10.3390/ijerph192013485>
9. Hofmann SG, Gómez AF. Mindfulness-based interventions for anxiety and depression. *Psychiatr Clin North Am* 2017; 40(4):739-749. DOI: <https://doi.org/10.1016/j.psc.2017.08.008>
10. Keng SL, Smoski MJ, Robins CJ. Effects of mindfulness on psychological health: A review of empirical studies. *Clin Psychol Rev* 2011; 31(6):1041-1056. DOI: <https://doi.org/10.1016/j.cpr.2011.04.006>
11. Parmentier FBR, García-Toro M, García-Campayo J, Yañez AM, Andrés P, Gili M. Mindfulness and symptoms of depression and anxiety in the general population: The mediating roles of worry, rumination, reappraisal and suppression. *Front Psychol* 2019; 10:506. DOI: <https://doi.org/10.3389/fpsyg.2019.00506>
12. Kachadourian LK, Harpaz-Rotem I, Tsai J, Southwick S, Pietrzak RH. Mindfulness as a mediator between trauma exposure and mental health outcomes: Results from the National Health and Resilience in Veterans Study. *Psychol Trauma* 2021; 13(2):223-230. DOI: <https://doi.org/10.1037/tra0000995>
13. Matiz A, Crescentini C, Bergamasco M, Budai R, Fabbro F. Inter-brain co-activations during mindfulness meditation. Implications for devotional and clinical settings. *Conscious Cogn* 2021; 95:103210. DOI: <https://doi.org/10.1016/j.concog.2021.103210>
14. Feruglio S, Matiz A, Pagnoni G, Fabbro F, Crescentini C. The impact of mindfulness meditation on the wandering mind: A systematic review. *Neurosci Biobehav Rev* 2021; 131:313-330. DOI: <https://doi.org/10.1016/j.neubiorev.2021.09.032>
15. McGee M. Meditation and psychiatry. *Psychiatry (Edgmont)* 2008; 5(1):28-41.

---

Received: June 11, 2024 | Revised: June 18, 2024 | Accepted: June 24, 2024

---